Super-PP-wave Algebra from Super-AdS×S Algebras in Eleven-dimensions

Machiko Hatsuda, Kiyoshi Kamimura[†] and Makoto Sakaguchi

Theory Division, High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki, 305-0801, Japan † Department of Physics, Toho University, Funabashi, 274-8510, Japan

E-mails: mhatsuda@post.kek.jp, kamimura@ph.sci.toho-u.ac.jp, Makoto.Sakaguchi@kek.jp

Abstract

Maximally supersymmetric spacetime algebras in eleven-dimensions, which are the isometry superalgebras of Minkowski space, $\operatorname{AdS}_7 \times S^4$, $\operatorname{AdS}_4 \times S^7$ and pp-wave background, are related by Inönü-Wigner contractions. The super- $\operatorname{AdS}_{4(7)} \times S^{7(4)}$ algebras allow to introduce two contraction parameters, the one for the flat limit to the super-Poincaré algebra and the other for a Penrose limit to the super-pp-wave algebra. Under these contractions supersymmetries are maintained because the Jacobi identity of three supercharges holds for any values of contraction parameters.

PACS: 11.30.Pb;11.17.+y;11.25.-w *Keywords:* Superalgebra; Anti-de Sitter; group contraction

1 Introduction

Recently, maximally supersymmetric pp-wave (plane-fronted gravitational wave with parallel rays) backgrounds of supergravity theories in eleven- and ten-dimensions have attracted great interests. In [1], maximally supersymmetric backgrounds in eleven-dimensions were classified into four types; flat Minkowski space (and their toroidal compactifications), $AdS_4 \times S^7$, $AdS_7 \times S^4$ and the super-pp-wave background. This super-pp-wave background in eleven-dimensions was reconsidered recently in [2]. For the type-IIB supergravity, a super-pp-wave background was discovered in [3].¹ The type-IIB superstrings in this background were shown to be exactly solvable [6] in spite of the presence of the RR five-form field strength. This model is expected to provide some hints for the study of superstrings on more general backgrounds [7][8][9].

The Penrose limit plays a central role in these recent studies of the pp-wave backgrounds. It was known that any solution of Einstein gravity admits plane-wave backgrounds in the Penrose limit [10]. This was extended to the solutions of supergravities in [11] and was studied in the context of the Wess-Zumino-Witten models in [12]. It was shown that the super-pp-wave background can be derived by the Penrose limit from the super-AdS×S backgrounds in [13] [14] and this was investigated further in [15]. In addition, the full supersymmetry algebra of the super-pp-wave background of the type-IIB theory was shown to be derived by an Inönü-Wigner (IW) contraction [16] from the super-AdS₅ × S^5 algebra of SU(2,2|4) in [17]. The Penrose limit was recognized to be important to explore the AdS/CFT correspondence [18] beyond massless string modes in [19]. This line was pursued further in [20].

In this paper, we consider the Penrose limit from an algebraic point of view, which is coordinate-independent and manifestly supersymmetric. We show that the isometry superalgebra of the super-pp-wave background in eleven-dimensions is obtained from the super-AdS₄₍₇₎ × $S^{7(4)}$ algebras by IW contractions. The supersymmetries are maintained during contractions because the Jacobi identity of three supercharges holds for any values of contraction parameters. In section 2, the pp-wave algebra is derived from the AdS₄₍₇₎ × $S^{7(4)}$ algebras. This is extended to the maximally supersymmetric case in section 3. The last section is devoted to a summary and discussions.

2 PP-wave Algebra from $AdS_{4(7)} \times S^{7(4)}$ Algebras

The $AdS_{p+2} \times S^{9-p}$ algebra, which is the isometry algebra of the $AdS_{p+2} \times S^{9-p}$ space, is given, in terms of dimensionless momenta P's and rotations J's, as

$$[P_{a}, P_{b}] = 4\epsilon^{2}J_{ab}, \qquad [P_{a'}, P_{b'}] = -\epsilon^{2}J_{a'b'}, [J_{ab}, P_{c}] = 2\eta_{bc}P_{a}, \qquad [J_{a'b'}, P_{c'}] = 2\eta_{b'c'}P_{a'}, J_{ab}, J_{cd}] = 4\eta_{ad}J_{bc}, \qquad [J_{a'b'}, J_{c'd'}] = 4\eta_{a'd'}J_{b'c'},$$
(2.1)

 $[[]J_{ab}, J_{cd}] = 4\eta_a$ ¹ For other dimensions see [4] and [5].

where $\epsilon^2 = 1$ for p = 2 and $\epsilon^2 = -1$ for p = 5. It is understood that the obvious antisymmetries of indices in the left hand side of equalities in (2.1) are to be implemented on the right hand side with unit weight. For p = 2, this algebra is the $AdS_4 \times S^7$ algebra with the vector index of AdS_4 , a = 0, 1, 2, 3, and that of S^7 , $a' = 4, ..., 9, \natural$. On the other hand, for p = 5, this algebra is the $AdS_7 \times S^4$ algebra with the vector index of S^4 , $a = \natural, 1, 2, 3$, and that of AdS_7 , a' = 4, ..., 9, 0. We use the metric $\eta_{\mu\nu}$ with $\eta_{00} = -1$ otherwise +1.

The symmetry group is $SO(p + 1, 2) \times SO(10 - p)$ which has a flat limit to $ISO(p + 1, 1) \times ISO(9 - p)$ by the following IW contraction

1. Rescale the translation generators P's as

$$P_a \to RP_a, \quad P_{a'} \to R'P_{a'}.$$
 (2.2)

where R/2 and R' are the radii of the $AdS_4(S^4)$ and that of $S^7(AdS_7)$, respectively.

2. Then take $R \to \infty$ and $R' \to \infty$.

In this limit, P's become the linear momenta and J's are the Lorentz generators. For the maximal spacetime supersymmetry, R and R' must satisfy

$$R = R', \quad R^2 = 2\rho R_S^2 = \frac{2}{\rho} R_{AdS}^2, \quad \rho = \begin{cases} 1/2 & \text{for } p = 2\\ 2 & \text{for } p = 5 \end{cases}$$
(2.3)

where R_S and R_{AdS} are the radii of S^{9-p} and that of AdS_{p+2} , respectively.

Besides the flat limit, as any other metrics, the $AdS \times S$ metric allows the Penrose limit [10] giving a plane wave metric. The Penrose limit can be understood as an IW contraction of the $AdS_{p+2} \times S^{9-p}$ algebra into the pp-wave algebra.

1. Define the light cone components of the momenta P's and *boost* generators P^* 's as

$$P_{\pm} \equiv \frac{1}{\sqrt{2}} (P_{\natural} \pm P_{0}), \quad P_{m} = (P_{i}, P_{i'}),$$

$$P_{m}^{*} = \begin{cases} (P_{i}^{*} = J_{i0}, P_{i'}^{*} = J_{i'\natural}) & \text{for } p = 2\\ (P_{i}^{*} = J_{i\natural}, P_{i'}^{*} = J_{i'0}) & \text{for } p = 5 \end{cases}$$
(2.4)

where i = 1, 2, 3, i' = 4, ..., 9, and m = (i, i').

2. Suppose the plane-wave propagates along x_+ time direction. The transverse translation and boost generators are rescaled with a dimensionless parameter Ω as

$$P_+ \to \frac{1}{\Omega^2} P_+ \quad P_m \to \frac{1}{\Omega} P_m \quad P_m^* \to \frac{1}{\Omega} P_m^*.$$
 (2.5)

3. Then take $\Omega \to 0$ limit.

To see them explicitly the $AdS_{p+2} \times S^{9-p}$ algebra (2.1) is rescaled, following to (2.2) and (2.5), as

$$\begin{split} [P_{i},P_{+}] &= 2\sqrt{2}\frac{\epsilon^{2}\Omega^{2}}{R^{2}}P_{i}^{*}, \qquad [P_{i'},P_{+}] = -\frac{\epsilon^{2}\Omega^{2}}{\sqrt{2}R^{2}}P_{i}^{*}, \\ [P_{i}^{*},P_{+}] &= -\frac{1}{\sqrt{2}}\epsilon^{2}\Omega^{2}P_{i}, \qquad [P_{i'}^{*},P_{+}] = \frac{1}{\sqrt{2}}\epsilon^{2}\Omega^{2}P_{i'}, \\ [P_{i},P_{-}] &= -2\sqrt{2}\frac{1}{R^{2}}P_{i}^{*}, \qquad [P_{i'},P_{-}] = -\frac{1}{\sqrt{2}R^{2}}P_{i'}^{*}, \\ [P_{i}^{*},P_{-}] &= \frac{1}{\sqrt{2}}P_{i}, \qquad [P_{i'}^{*},P_{-}] = \frac{1}{\sqrt{2}}P_{i'}, \\ [P_{i}^{*},P_{-}] &= \frac{1}{\sqrt{2}}\eta_{ij}(\epsilon^{2}\Omega^{2}P_{-}-P_{+}), \qquad [P_{i'}^{*},P_{j'}] = -\frac{1}{\sqrt{2}}\eta_{i'j'}(\epsilon^{2}\Omega^{2}P_{-}+P_{+}), \\ [P_{i},P_{j}] &= 4\frac{\epsilon^{2}\Omega^{2}}{R^{2}}J_{ij}, \qquad [P_{i'},P_{j'}] = -\frac{\epsilon^{2}\Omega^{2}}{R^{2}}J_{i'j'}, \\ [P_{i}^{*},P_{j}^{*}] &= \epsilon^{2}\Omega^{2}J_{ij}, \qquad [P_{i'}^{*},P_{j'}] = -\epsilon^{2}\Omega^{2}J_{i'j'}, \\ [J_{ij},P_{k}] &= 2\eta_{jk}P_{i}, \qquad [J_{i'j'},P_{k'}] = 2\eta_{j'k'}P_{i'}, \\ [J_{ij},J_{kl}] &= 4\eta_{il}J_{jk}, \qquad [J_{i'j'},P_{k'}] = 2\eta_{j'k'}P_{i'}^{*}, \\ [J_{ij},J_{kl}] &= 4\eta_{il}J_{jk}, \qquad [J_{i'j'},J_{k'l'}] = 4\eta_{i'l'}J_{j'k'}, \end{aligned}$$

$$(2.6)$$

where we used relation R = R' of (2.3) for simplicity. By taking $\Omega \to 0$ limit, the ϵ dependence, which distinguishes the p = 2 case from the p = 5 case, disappears and (2.6) becomes the unique plane wave algebra in eleven-dimensions,

$$[P_{i}, P_{-}] = -2\sqrt{2} \frac{1}{R^{2}} P_{i}^{*}, \qquad [P_{i'}, P_{-}] = -\frac{1}{\sqrt{2}R^{2}} P_{i'}^{*},$$

$$[P_{m}^{*}, P_{-}] = \frac{1}{\sqrt{2}} P_{m}, \qquad [P_{m}^{*}, P_{n}] = -\frac{1}{\sqrt{2}} \eta_{mn} P_{+},$$

$$[J_{mn}, P_{p}] = 2\eta_{np} P_{m}, \qquad [J_{mn}, P_{p}^{*}] = 2\eta_{np} P_{m}^{*},$$

$$[J_{mn}, J_{pq}] = 4\eta_{mq} J_{np}.$$
(2.7)

This is the symmetry algebra of the pp-wave metric [21]. The Poincaré algebra of ISO(10, 1) group is obtained by taking flat limit $R \to 0$ in (2.7) and by making manifest restored symmetry generators such as $J_{ij'}$, $J_{\pm i}$, etc.

3 Super-PP-wave algebra from Super-AdS $_{4(7)} \times S^{7(4)}$ Algebras

We extend the previous analysis to the supersymmetric case and show the maximally supersymmetric eleven-dimensional pp-wave algebra [2] is obtained by IW contractions of the super-AdS_{p+2} × S^{9-p} algebras. In this case the Jacobi identities of the superalgebra give a restriction on the radii of AdS and S, R_{AdS} and R_S , as $R_{AdS} = \frac{1}{2}R_S$ for p = 2 and $R_{AdS} = 2R_S$ for p = 5, as in (2.3). In eleven-dimensions, the supersymmetry generators Q_{α} are 32 Majorana spinors. The gamma matrices Γ^{μ} , $(\mu = 0, 1, ..., 9, \natural)$, in eleven-dimensions are composed in terms of those of AdS₄, γ^a , (a = 0, 1, 2, 3), and of S^7 , $\gamma^{a'}$, $(a' = 4, 5, ..., 9, \natural)$, or in terms of those of S^4 , γ^a , $(a = \natural, 1, 2, 3)$, and of AdS₇, $\gamma^{a'}$, (a' = 4, 5, ..., 9, 0), as ²

$$\Gamma^{a} = \gamma^{a} \otimes \mathbb{1}, \quad \Gamma^{a'} = \gamma_{5} \otimes \gamma^{a'}, \quad \gamma_{5} = \begin{cases} i\gamma^{0123} & \text{for } p = 2\\ \gamma^{\natural 123} & \text{for } p = 5 \end{cases}.$$
(3.1)

They satisfy $\{\Gamma^{\mu}, \Gamma^{\nu}\} = 2\eta^{\mu\nu}$. The charge conjugation matrix \mathcal{C} in eleven-dimensions is taken as

$$\mathcal{C} = C \otimes C' \tag{3.2}$$

where C and C' are the charge conjugation matrices for AdS_4 (S⁴) and S⁷ (AdS₇) spinors respectively.

The bosonic part of the super-AdS_{p+2} × S^{9-p} algebra is (2.1). In addition to it, the odd generators Q_{α} satisfy

$$[P_{a},Q] = i\epsilon Q\gamma_{5}\gamma_{a} \otimes \mathbb{1}, \qquad [P_{a'},Q] = \frac{i}{2}\epsilon Q\mathbb{1} \otimes \gamma_{a'},$$

$$[J_{ab},Q] = \frac{1}{2}Q\gamma_{ab} \otimes \mathbb{1}, \qquad [J_{a'b'},Q] = \frac{1}{2}Q\mathbb{1} \otimes \gamma_{a'b'},$$

$$\{Q,Q\} = -2C\gamma^{a} \otimes C'P_{a} - 2C\gamma_{5} \otimes C'\gamma^{a'}P_{a'}$$

$$-2i\epsilon C\gamma_{5}\gamma^{ab} \otimes C'J_{ab} + i\epsilon C\gamma_{5} \otimes C'\gamma^{a'b'}J_{a'b'}.$$
(3.3)

The Jacobi identities are satisfied as long as $\epsilon^2 = 1$ for p = 2 and $\epsilon^2 = -1$ for p = 5. Hereafter, we choose $\epsilon = 1$ for p = 2 and $\epsilon = -i$ for p = 5, for presentation.

For the present purpose, we rewrite this superalgebra in terms of eleven-dimensional covariant gamma matrices, Γ 's, instead of γ 's. It is convenient to introduce following matrix

$$I = \Gamma^{123}, \quad \Gamma^{\sharp 123} = \Gamma^{\sharp} I = -i\epsilon\gamma_5 \otimes \mathbb{1}, \quad \Gamma^{\sharp} = \begin{cases} \Gamma^0 & \text{for } p = 2\\ -\Gamma^{\natural} & \text{for } p = 5 \end{cases}.$$
(3.4)

² For the gamma-matrices in four-dimensions, we use a Majorana representation for AdS_4 (a = 0, 1, 2, 3) and a pseudo-symplectic representation for S^4 ($a = \natural, 1, 2, 3$) which satisfy, in both cases,

$$\tilde{C} = -C, \quad \tilde{\gamma}^a = -C\gamma^a C^{-1}, \quad \{\gamma^a, \gamma^b\} = 2\eta^{ab}$$

and $C\gamma^{a_1\cdots a_N}$ is symmetric iff $N = 1, 2 \mod 4$. For the gamma-matrices in seven-dimensions, we use a pseudo-Majorana representation for S^7 $(a' = 4, 5, ..., 9, \natural)$ and a symplectic Majorana representation for AdS₇ $(a' = 4, 5, ..., 9, \natural)$ which satisfy, in both cases,

$$\tilde{C}' = +C', \quad \tilde{\gamma}^{a'} = -C'\gamma^{a'}C'^{-1}, \quad \{\gamma^{a'}, \gamma^{b'}\} = 2\eta^{a'b'}$$

and $C'\gamma^{a'_1\cdots a'_N}$ is symmetric iff $N=0,3 \mod 4$.

The commutation relations (3.3) for the super-AdS_{p+2} × S^{9-p} algebra are rewritten as

$$[P_a, Q] = -Q\Gamma^{\sharp}I\Gamma_a, \quad [P_{a'}, Q] = -\frac{1}{2}Q\Gamma^{\sharp}I\Gamma_{a'}, \quad [J_{mn}, Q] = \frac{1}{2}Q\Gamma_{mn}$$

$$\{Q, Q\} = -2\mathcal{C}\Gamma^{\mu}P_{\mu} + 2\mathcal{C}\Gamma^{\sharp}I\Gamma^{ab}J_{ab} - \mathcal{C}\Gamma^{\sharp}I\Gamma^{a'b'}J_{a'b'}.$$
(3.5)

Associating with the rescaling of P's by R as in (2.2) with (2.3), the dimensionless supercharges Q_{α} are rescaled as

$$P_a \to RP_a \ , \ P_{a'} \to RP_{a'} \ , \ Q_\alpha \to \sqrt{R} \ Q_\alpha.$$
 (3.6)

Corresponding to the rescaling of bosonic generators (2.5), the components of the supercharges must be rescaled with proper weights. To do this, we decompose supercharges Qas

$$Q = Q_{+} + Q_{-}, \quad Q_{\pm} = Q_{\pm} \mathcal{P}_{\pm},$$
 (3.7)

using the light cone projection operators

$$\mathcal{P}_{\pm} = \frac{1}{2} \Gamma_{\pm} \Gamma_{\mp}, \quad \Gamma_{\pm} \equiv \frac{1}{\sqrt{2}} (\Gamma_{\natural} \pm \Gamma_0). \tag{3.8}$$

In order to obtain the well defined limit of the super-AdS_{p+2} × S^{9-p} algebra, the supercharges turn out to be rescaled as

$$Q_+ \to \frac{1}{\Omega} Q_+ \quad , \quad Q_- \to Q_-.$$
 (3.9)

The super-AdS_{p+2} × S^{9-p} algebra (3.5), after the rescaling (3.6) by R and (2.5) and (3.9) by Ω , becomes

$$\begin{split} [P_{+},Q_{+}] &= \frac{\Omega^{2}\epsilon^{2}}{2\sqrt{2R}}Q_{+}I, \qquad [P_{-},Q_{+}] = -\frac{3}{2\sqrt{2R}}Q_{+}I, \\ [P_{i},Q_{-}] &= \frac{1}{\sqrt{2R}}Q_{+}\Gamma^{-}I\Gamma_{i}, \qquad [P_{i'},Q_{-}] = \frac{1}{2\sqrt{2R}}Q_{+}\Gamma^{-}I\Gamma_{i'}, \\ [P_{m}^{*},Q_{-}] &= \frac{1}{2\sqrt{2}}Q_{+}\Gamma_{m}\Gamma^{-}, \qquad [J_{mn},Q_{\pm}] = \frac{1}{2}Q_{\pm}\Gamma_{mn} \\ [P_{+},Q_{-}] &= \frac{3\Omega^{2}\epsilon^{2}}{2\sqrt{2R}}Q_{-}I, \qquad [P_{-},Q_{-}] = -\frac{1}{2\sqrt{2R}}Q_{-}I, \\ [P_{i},Q_{+}] &= -\frac{\Omega^{2}\epsilon^{2}}{\sqrt{2R}}Q_{-}\Gamma^{+}I\Gamma_{i}, \qquad [P_{i'},Q_{+}] = -\frac{\Omega^{2}\epsilon^{2}}{2\sqrt{2R}}Q_{-}\Gamma^{+}I\Gamma_{i'}, \\ [P_{i}^{*},Q_{+}] &= -\frac{\Omega^{2}\epsilon^{2}}{2\sqrt{2}}Q_{-}\Gamma_{i}\Gamma^{+}, \qquad [P_{i'}^{*},Q_{+}] = \frac{\Omega^{2}\epsilon^{2}}{2\sqrt{2}}Q_{-}\Gamma_{i'}\Gamma^{+}, \\ \{Q_{+},Q_{+}\} &= -2\mathcal{C}\Gamma^{+}P_{+} + \frac{\sqrt{2}\Omega^{2}\epsilon^{2}}{R}\mathcal{C}\Gamma^{+}I\Gamma^{ij}J_{ij} - \frac{\Omega^{2}\epsilon^{2}}{\sqrt{2R}}\mathcal{C}\Gamma^{+}I\Gamma^{i'j'}J_{i'j'}, \\ \{Q_{-},Q_{-}\} &= -2\mathcal{C}\Gamma^{-}P_{-} - \frac{\sqrt{2}}{R}\mathcal{C}\Gamma^{-}I\Gamma^{ij}J_{ij} + \frac{1}{\sqrt{2}\Omega}\mathcal{C}\Gamma^{-}I\Gamma^{i'j'}J_{i'j'}, \end{split}$$

$$\{Q_+, Q_-\} = \left(-2\mathcal{C}\Gamma^m P_m - \frac{4}{R}\mathcal{C}I\Gamma^i P_i^* - \frac{2}{R}\mathcal{C}I\Gamma^{i'} P_{i'}^*\right)\mathcal{P}_-.$$
(3.10)

It is important that negative power terms of Ω disappear owing to the presence of the light cone projections. Therefore we can take the consistent Penrose limit $\Omega \to 0$ of the algebra to obtain

$$\begin{split} [P_{-},Q_{+}] &= -\frac{3}{2\sqrt{2R}}Q_{+}I, \qquad [P_{-},Q_{-}] = -\frac{1}{2\sqrt{2R}}Q_{-}I, \\ [P_{i},Q_{-}] &= \frac{1}{\sqrt{2R}}Q_{+}\Gamma^{-}I\Gamma_{i}, \qquad [P_{i'},Q_{-}] = \frac{1}{2\sqrt{2R}}Q_{+}\Gamma^{-}I\Gamma_{i'}, \\ [P_{m}^{*},Q_{-}] &= \frac{1}{2\sqrt{2}}Q_{+}\Gamma_{m}\Gamma^{-}, \qquad [J_{mn},Q_{\pm}] = \frac{1}{2}Q_{\pm}\Gamma_{mn}, \end{split}$$

$$\{Q_{+}, Q_{+}\} = -2\mathcal{C}\Gamma^{+}P_{+}$$

$$\{Q_{-}, Q_{-}\} = -2\mathcal{C}\Gamma^{-}P_{-} - \frac{\sqrt{2}}{R}\mathcal{C}\Gamma^{-}I\Gamma^{ij}J_{ij} + \frac{1}{\sqrt{2R}}\mathcal{C}\Gamma^{-}I\Gamma^{i'j'}J_{i'j'},$$

$$\{Q_{+}, Q_{-}\} = \left(-2\mathcal{C}\Gamma^{m}P_{m} - \frac{4}{R}\mathcal{C}I\Gamma^{i}P_{i}^{*} - \frac{2}{R}\mathcal{C}I\Gamma^{i'}P_{i'}^{*}\right)\mathcal{P}_{-}.$$

$$(3.11)$$

After the Penrose limit, all ϵ dependence have disappeared. This reflects the fact that the super-AdS₄ × S⁷ background and the super-AdS₇ × S⁴ background reduce into the unique super-pp-wave background in the Penrose limit. In this way, we have obtained the super-pp-wave algebra from the super-AdS₄ × S⁷ algebra and the super-AdS₇ × S⁴ algebra.

Furthermore the flat limit to super-Poincaré algebra can be taken by $R \to \infty$ in (2.7) and (3.11), and by making manifest restored (bosonic) symmetry generators as was mentioned in the last section.

We have established relations of the maximally supersymmetric spacetime algebras in eleven-dimensions; super-Poincaré, super-AdS₄ × S^7 , super-AdS₇ × S^4 and super-pp-wave, by IW contractions of superalgebras.

4 Summary and Discussions

We derived the super-pp-wave algebra form the super-AdS₄₍₇₎ × $S^{7(4)}$ algebras in elevendimensions by IW contractions which correspond to the Penrose limits of the super-AdS₄₍₇₎ × $S^{7(4)}$ backgrounds. The differences between the super-AdS₄ × S^7 algebra and the super-AdS₇× S^4 algebra, which are essentially caused from interchanging 0-component and \natural -component, are shown to disappear after the contraction resulting to the unique super-pp-wave algebra in eleven-dimensions. A solution of plane wave is a function of $x_+ = t + x_{\natural}$ and it is invariant under interchanging t and x_{\natural} . The Penrose limit brings to such a plane wave space. This naturally explains why the maximally supersymmetric pp-wave algebra is unique in spite of the presence of two distinct super-AdS algebras in eleven-dimensions. This property will be deeply related to the T-duality discussed in [11]. Supersymmetries are maintained during contractions because the Jacobi identities hold for any values of contraction parameters, R and Ω , and even for their limits $R \to \infty$ and $\Omega \to 0$.

It is noted that one can relate the super-pp-algebra, (2.7) and (3.11), to the one obtained in [2]. To do this, we rewrite generators as

$$P_{\pm} \to 3\sqrt{2}e_{\pm}, \quad P_m \to 3\sqrt{2}e_m, \\ P_i^* \to \frac{9\sqrt{2}}{2\mu^2}e_i^*, \quad P_{i'}^* \to \frac{18\sqrt{2}}{\mu^2}e_{i'}^*, \quad Q_{\pm} \to \sqrt{6\sqrt{2}}Q_{\pm}$$
(4.12)

where $\mu = 1/R$. Under this, the super-pp-wave algebra (2.7) and (3.11) turns out to be

$$[e_{m}, e_{-}] = -e_{m}^{*}, \quad [e_{i}^{*}, e_{-}] = \frac{\mu^{2}}{9}e_{i}, \quad [e_{i'}^{*}, e_{-}] = \frac{\mu^{2}}{36}e_{i'},$$

$$[e_{i}^{*}, e_{j}] = -\frac{\mu^{2}}{9}\eta_{ij}e_{+}, \quad [e_{i'}^{*}, e_{j'}] = -\frac{\mu^{2}}{36}\eta_{i'j'}e_{+},$$

$$[J_{mn}, e_{p}] = 2\eta_{np}e_{m}, \quad [J_{mn}, e_{p}^{*}] = 2\eta_{np}e_{m}^{*}, \quad [J_{mn}, J_{pq}] = 4\eta_{np}J_{mq},$$

$$[e_{-}, Q_{+}] = -\frac{\mu}{4}Q_{+}I, \quad [e_{-}, Q_{-}] = -\frac{\mu}{12}Q_{-}I, \quad [e_{i}, Q_{-}] = \frac{\mu}{6}Q_{+}\Gamma^{-}I\Gamma_{i},$$

$$[e_{i'}, Q_{-}] = \frac{\mu}{12}Q_{+}\Gamma^{-}I\Gamma_{i'}, \quad [e_{i}^{*}, Q_{-}] = \frac{\mu^{2}}{18}Q_{+}\Gamma_{i}\Gamma^{-}, \quad [e_{i'}^{*}, Q_{-}] = \frac{\mu^{2}}{72}Q_{+}\Gamma_{i'}\Gamma^{-},$$

$$\{Q_{+}, Q_{+}\} = -\mathcal{C}\Gamma^{+}e_{+},$$

$$\{Q_{-}, Q_{-}\} = -\mathcal{C}\Gamma^{-}e_{-} - \frac{\mu}{6}\mathcal{C}\Gamma^{-}I\Gamma^{ij}J_{ij} + \frac{\mu}{12}\mathcal{C}\Gamma^{-}I\Gamma^{i'j'}J_{i'j'},$$

$$\{Q_{+}, Q_{-}\} = -\mathcal{C}\Gamma^{m}e_{m} - \frac{3}{\mu}\mathcal{C}I\Gamma^{i}e_{i}^{*} - \frac{6}{\mu}\mathcal{C}I\Gamma^{i'}e_{i'}^{*}.$$

$$(4.13)$$

and is the superalgebra in [2] ³. Our presentation reveals the way to take the flat limit while the flat limit $\mu \to 0$ is not obvious in the form of (4.13).

The relation between the super-pp wave algebra and the $AdS_{4(7)} \times S^{7(4)}$ algebra is useful for constructing mechanical actions of M-branes in the super-pp wave background. Any form field in the former can be derived from the corresponding form in the latter. This approach will make the (super)symmetry of the pp-wave systems manifest.

References

- J. Kowalski-Glikman, "Vacuum states in supersymmetric Kaluza-Klein theory," Phys. Lett. B 134 (1984) 194; P. T. Chrusciel and J. Kowalski-Glikman, "The Isometry Group And Killing Spinors For The P P Wave Space-Time In D = 11 Supergravity," Phys. Lett. B 149 (1984) 107.
- [2] J. Figueroa-O'Farrill and G. Papadopoulos, "Homogeneous fluxes, branes and a maximally supersymmetric solution of M-theory", JHEP 0108 (2001) 036, hep-th/0105308.

³ In the algebra given in [2], Q_+ and Q_- are exchanged in the commutation relations of $[e_m^*, Q_-]$ and $[e_m, Q_-]$. It apparently breaks the Ω -grading property which played the central role in the Penrose limit as explained in this paper.

- [3] M. Blau, J. Figueroa-O'Farrill, C. Hull and G. Papadopoulos, "A new maximally supersymmetric background of IIB superstring theory," JHEP **0201** (2002) 047, hep-th/0110242.
- [4] J. Kowalski-Glikman, "Positive Energy Theorem And Vacuum States For The Einstein-Maxwell System," Phys. Lett. B 150 (1985) 125.
- [5] P. Meessen, "A small note on PP-wave vacua in 6 and 5 dimensions," Phys. Rev. D 65 (2002) 087501, hep-th/0111031.
- [6] R. R. Metsaev, "Type IIB Green-Schwarz superstring in plane wave Ramond-Ramond background," Nucl. Phys. B 625 (2002) 70, hep-th/0112044.
- [7] R. R. Metsaev and A. A. Tseytlin, "Exactly solvable model of superstring in plane wave Ramond-Ramond background," hep-th/0202109; J.G. Russo and A.A. Tseytlin, "On solvable models of type IIB superstring in NS-NS and R-R plane wave backgrounds", hepth/0202179.
- [8] M. Billó and I. Pesando, "Boundary states for GS superstrings in an Hpp wave background", hep-th/0203028; J. Michelson, "(Twisted) Toroidal compactification of pp-waves", hepth/0203140.
- [9] N. Berkovits, "Conformal field theory for the superstring in a Ramond-Ramond plane wave background", hep-th/0203248.
- [10] R. Penrose, "Any space-time has a plane wave as a limit", in Differential Geometry and Relativity, Cahen and Flato eds. (1976) Reidel Publishing, Dordrecht-Holland.
- [11] R. Güven, "Plane wave limits and T-duality", Phys.Lett. B482 (2000) 255-263, hepth/0005061.
- [12] K. Sfetsos, "Gauging a nonsemisimple WZW model", Phys. Lett. B324 (1994) 335, hep-th/9311010; D.I. Olive, E. Rabinovici and A. Schwimmer, "A class of string backgrounds as a semiclassical limit of WZW models", Phys. Lett. B321 (1994) 361, hepth/9311081; K. Sfetsos and A.A. Tseytlin, "Four-dimensional plane wave string solutions with coset CFT description", Nucl. Phys. B427 (1994) 245, hep-th/9404063; A.A. Tseytlin, "Exact string solutions and duality", hep-th/9407099.
- [13] M. Blau, J. Figueroa-O'Farrill, C. Hull and G. Papadopoulos, "Penrose limits and maximal supersymmetry," hep-th/0201081.
- [14] M. Blau, J. Figueroa-O'Farrill and G. Papadopoulos, "Penrose limits, supergravity and brane dynamics," hep-th/0202111.
- [15] M. Alishahiha and M.M. Sheikh-Jabbari, "The pp-wave limits of orbifolded AdS₅ × S⁵" hep-th/0203018; M. Cvetič, H. Lü and C.N. Pope, "Penrose limits, pp-waves and deformed M2-branes", hep-th/0203082; U. Gürsoy, C. Núñez and M. Schvellinger, "RG flows from Spin(7), CY 4-fold and HK manifolds to AdS, penrose limits and pp waves", hep-th/0203124; E. Floratos and A. Kehagias, "Penrose limits of orbifolds and orientifolds", hep-th/0203134; R. Güven, "Randall-Sundrum zero mode as a Penrose limit", hep-th/0203153; M. Cvetič, H. Lü and C.N. Pope, "M-theory pp-waves, Penrose limits and supernumerary supersymmetries", hep-th/0203229; J.P. Gauntlett and C.M. Hull, "pp-waves in 11-dimensions with extra supersymmetry", hep-th/0203255.

- [16] E. Inönü and E. P. Wigner, "On the contraction of groups and their representations," Proc. Nat. Acad. Sci. **39** (1953) 510; E. Inönü, "Contraction of Lie group and their representations", in Group Theoretical Concepts and Methods in Elementary Particle Physics, ed. F.Gürsey (Gordon and Breach, New York, 1964).
- [17] M. Hatsuda, K. Kamimura and M. Sakaguchi, "From super-AdS₅ × S^5 algebra to super-pp-wave algebra", hep-th/0202190, to appear in NPB.
- [18] J. Maldacena, "The large N limit of superconformal field theories and supergravity," Adv. Theor. Math. Phys. 2 (1998) 231, Int. J. Theor. Phys. 38 (1998) 1113, hep-th/9711200; O. Aharony, S. S. Gubser, J. Maldacena, H. Ooguri and Y. Oz, "Large N field theories, string theory and gravity," Phys. Rept. 323 (2000) 183, hep-th/9905111.
- [19] D. Berenstein, J. Maldacena and H. Nastase, "Strings in flat space and pp waves from $\mathcal{N} = 4$ super Yang Mills", hep-th/0202021.
- [20] N. Itzhaki, I.R. Klebanov and S. Mukhi, "PP wave limit and enhanced supersymmetry in gauge theories", hep-th/0202153; J. Gomis and H. Ooguri, "Penrose limit of N=1 gauge theories", hep-th/0202157; L.A. Pando Zayas and J. Sonnenschein, "On Penrose limits and gauge theories", hep-th/0202186; N. Kim, A. Pankiewicz, S.-J. Rey and S. Theisen, "Superstring on pp-wave orbifold from large-N quiver gauge theory", hep-th/0203080; T. Takayanagi and S. Terashima, "Strings on orbifolded pp-waves", hep-th/0203093; S.R. Das, C. Gomez, S.-J. Rey, "Penrose limit, spontaneous symmetry breaking and holography in pp-wave background", hep-th/0203164; A. Dabholkar and S. Parvizi, "Dp branes in pp-wave background", hep-th/0203231; D. Berenstein, E. Gava, J. Maldacena, K.S. Narain and H. Nastase, "Open strings on plane waves and their Yang-Mills duals", hep-th/0203249; P. Lee and J. Park, "Open strings in pp-wave background from defect conformal field theory", hep-th/0203257.
- [21] D. Kramer, H. Stephani, E. Herlt and M. MacCallum, "The plane-fronted gravitational waves with parallel rays (pp waves)", p.233-236, in Exact solutions of Einstein's field equation (1980), Cambridge University Press, Cambridge.